

- 5 Kim RJ, Fieno DS, Parrish TB, *et al.* Relationship of MRI delayed contrast enhancement to irreversible injury, infarct age, and contractile function. *Circulation* 1999;100:1992–2002.
- ▶ **Landmark study comparing contrast enhancement to histopathology and concluding that contrast enhancement occurs in the setting of irreversible myocardial injury only.**
- 6 Ansari M, Araoz PA, Gerard SK, *et al.* Comparison of late enhancement cardiovascular magnetic resonance and thallium SPECT in patients with coronary disease and left ventricular dysfunction. *J Cardiovasc Magn Reson* 2004;6:549–56.
- 7 Klein C, Nekolla SG, Bengel FM, *et al.* Assessment of myocardial viability with contrast-enhanced magnetic resonance imaging: comparison with positron emission tomography. *Circulation* 2002;105:162–7.
- 8 Kühl HP, Beek AM, van der Weerd AP, *et al.* Myocardial viability in chronic ischemic heart disease: comparison of contrast-enhanced magnetic resonance imaging with (18)F-fluorodeoxyglucose positron emission tomography. *J Am Coll Cardiol* 2003;41:1341–8.
- 9 Wagner A, Mahrholdt H, Holly TA, *et al.* Contrast-enhanced MRI and routine single photon emission computed tomography (SPECT) perfusion imaging for detection of subendocardial myocardial infarcts: an imaging study. *Lancet* 2003;361:374–9.
- ▶ **Landmark study concluding that ceCMR detects myocardial infarcts which are missed by SPECT imaging.**
- 10 Mahrholdt H, Wagner A, Parker M, *et al.* Relationship of contractile function to transmural extent of infarction in patients with chronic coronary artery disease. *J Am Coll Cardiol* 2003;42:505–12.
- 11 Hillenbrand HB, Kim RJ, Parker MA, *et al.* Early assessment of myocardial salvage by contrast-enhanced magnetic resonance imaging. *Circulation* 2000;102:1678–83.
- 12 Choi K, Kim RJ, Gubernikoff G, *et al.* The transmural extent of acute myocardial infarction predicts long term improvement in contractile function. *Circulation* 2001;104:1101–7.
- 13 Kim RJ, Wu E, Rafael A, *et al.* The use of contrast-enhanced magnetic resonance imaging to identify reversible myocardial dysfunction. *N Engl J Med* 2000;343:1445–53.
- ▶ **The first study to demonstrate that ceCMR can accurately predict reversible myocardial dysfunction in patients with ischaemic heart disease before coronary revascularisation.**
- 14 Wellnhofer E, Olariu A, Klein C, *et al.* Magnetic resonance low-dose dobutamine test is superior to scar quantification for the prediction of functional recovery. *Circulation* 2004;109:2172–4.
- 15 Nagel E, Lehmkühl HB, Bocksch W, *et al.* Diagnosis of ischemia-induced wall motion abnormalities with the use of high-dose dobutamine stress MRI: comparison with dobutamine stress echocardiography. *Circulation* 1999;99:763–70.
- ▶ **Excellent study demonstrating that the diagnostic performance of dobutamine CMR is at least comparable to dobutamine echocardiography and superior in patients with poor acoustic windows.**
- 16 Hundley WG, Hamilton CA, Thomas MS, *et al.* Utility of fast cine magnetic resonance imaging and display for the detection of myocardial ischemia in patients not well suited for second harmonic stress echocardiography. *Circulation* 1999;100:1676–9.
- 17 Klocke FJ, Simonetti OP, Judd RM, *et al.* Limits of detection of regional differences in vasodilated flow in viable myocardium by first-pass magnetic resonance perfusion imaging. *Circulation* 2001;104:2412–16.
- 18 Schwitler J, Nanz D, Kneifel S, *et al.* Assessment of myocardial perfusion in coronary artery disease by magnetic resonance: a comparison with positron emission tomography and coronary angiography. *Circulation* 2001;103:2230–35.
- 19 Klem I, Heitner JF, Shah DJ, *et al.* Improved detection of coronary artery disease by stress perfusion cardiovascular magnetic resonance with the use of delayed enhancement infarction imaging. *J Am Coll Cardiol* 2006;47:1630–8.
- ▶ **The first study evaluating an algorithmic approach using perfusion CMR as well as ceCMR for the initial diagnosis of CAD.**
- 20 Paetsch I, Jahnke C, Wahl A, *et al.* Comparison of dobutamine stress magnetic resonance, adenosine stress magnetic resonance, and adenosine stress magnetic resonance perfusion. *Circulation* 2004;110:835–42.

MULTIPLE CHOICE QUESTIONS

Education in Heart Interactive (www.heartjnl.com/misc/education.shtml)

There are six multiple choice questions associated with each *Education in Heart* article (these questions have been written by the authors of the articles). Each article is submitted to EBAC (European Board for Accreditation in Cardiology; www.ebac-cme.org) for 1 hour of external CPD credit.

How to find the MCQs: Click on the **Online Learning**: [[Take interactive course](#)] link on the table of contents for the issue online or on the *Education in Heart* collection (www.heartjnl.com/cgi/collection/heart_education).

Free access: This link will take you to the BMJ Publishing Group's online learning website. Your *Heart Online* user name and password will be recognised by this website. As a *Heart* subscriber you have free access to these MCQs but you must register on the site so you can track your learning activity and receive credit for completed courses.

How to get access: If you have not yet activated your *Heart Online* access, please do so by visiting <http://www.bmjournals.com/cgi/activate/basic> and entering your six digit (all numeric) customer number (found above your address label with your print copy). If you have any trouble activating or using the site please contact subscriptions@bmjgroup.com

Case based Heart: You might also be interested in the interactive cases published in association with *Heart* (http://cpd.bmjournals.com/cgi/hierarchy/cpd_node;CBH)